

AGRORAM 16 COMPLEX




PREVENTS AND TREATS COPPER DEFICIENCY ALLOWED IN ORGANIC FARMING

The copper present in AGRORAM 16 COMPLEX is bound to a particular organic complex that enhances its activity, favoring its absorption. Moreover the presence of other oligoelements increases its efficacy and makes it particularly useful in solving very common deficiency conditions during the first vegetative phases.

CROP	TIME OF APPLICATION	DOSE/HECTARE*
Kiwifruit, Stone fruits (Apricot, Cherry, Nectarine, Peach, Plum), Citrus (Orange, Bergamot, Clementine, Lemon, Tangerine), Pome fruits (Quince, Apple, Pear), Olive e Grapes	At early vegetative phases, at pre-flowering or at the appearance of the first deficiency symptoms	1-2 Kg
Fruiting vegetables (Watermelon, Cucumber, Eggplant, Melon, Pepper, Tomato, Zucchini, Pumpkin)	At early vegetative phases, at pre-flowering or at the appearance of the first deficiency symptoms. To be repeated 2-3 times every 8-10 days	1-2 Kg
Cereal crops (Oats, Wheat, Corn, Barley, Rice, Rye, Sorghum, Triticale)	At early vegetative phases, at pre-flowering or at the appearance of the first deficiency symptoms	1-2 Kg
Beets e Tobacco	At early vegetative phases, at pre-flowering or at the appearance of the first deficiency symptoms. To be repeated 2-3 times every 8-10 days	1-2 Kg
Industrial crops (Sugarcane, Rapeseed, Cotton, Sunflower, Industrial tomato, Soybeans)	At early vegetative phases, at pre-flowering or at the appearance of the first deficiency symptoms	1-2 Kg
Flower crops (geranium and chrysanthemum in particular)	At early vegetative phases, at pre-flowering or at the appearance of the first deficiency symptoms. To be repeated 2-3 times every 8-10 days	1-2 Kg

COMPOSITION	
Copper (Cu) soluble in water	16.00%
Manganese (Mn) soluble in water	2.00%
Molybdenum (Mo) soluble in water	0.02%
Zinc (Zn) soluble in water	1.00%

PHYSICO-CHEMICAL FEATURES	
SOLUBLE POWDER	
pH (sol 1%)	4.0
Conductivity E.C. $\mu\text{S}/\text{cm}$ (1‰)	1060
METHOD OF USE	
	Foliar fertilization

PACKAGING: 5 - 25 KG

The choice of the dose is subordinated to various factors and can be varied when necessary. All applications can be repeated in relation to the different crop needs. You can contact our Technical Service for the correct application on specific soils and under specific climate conditions.