

## IMPROVES THE RATE OF CHLOROPHYLL BIOSYNTHESIS AND PHOTOSYNTHESIS

## IMPROVES TOLERANCE TO ADVERSE ENVIRONMENTAL CONDITIONS


## IMPROVES THE PLANT'S NUTRIENT UTILIZATION EFFICIENCY

SKICC is the biostimulant specifically conceived to promote chlorophyll biosynthesis and thus improve photosynthesis and plants' metabolism ensuring the reaching of optimal crops' yield. ADRIATICA conceived SKICC by employing an internally developed manufacturing process (KK Technology) aimed at making nutrients' uptake faster and more efficient. Its formulation has a balanced nutritional ratio that, in addition to improving chlorophyll synthesis, optimizes nutritional uptake and its use efficiency. All this makes SKICC highly efficient in terms of produce quality and yield.

CE BIOSTIMULANT CERTIFIED BY AGENCY CODE. NANDO 2832 WITH N° CERT.: DE.4161

CROP	TIME OF APPLICATION	DOSE/HECTARE*
Fruiting vegetables (Pumpkin, Zucchini, Tomato, Pepper, Melon, Eggplant, Cucumber, Watermelon)	Post-transplanting, pre-flowering, veraison (change of color), combined with phytosanitary treatments	4-6 kg
Leafy vegetables (Spinach, Celery, Escarole, Rocket, Radicchio, Lettuce, Chicory)	Post-transplanting and during vegetative development combined with phytosanitary treatments	4-6 kg
Other vegetables (Leek, Fennel, Onion, Cauliflower, Cabbage, Broccoli, Garlic)	Post-transplanting and during vegetative development combined with phytosanitary treatments	4-6 kg
Corn e Sunflower	3-6 leaf combined with post-emergence herbicide treatment	6-8 kg
Cereal crops (Wheat, Barley, Rice)	Stem elongation combined with post-emergence herbicide treatment, booting phase combined with fungicide treatment	4-6 kg
Rapeseed	Pre-flowering and fruit set combined with insecticide treatment	4-6 kg
Beets	Leaf canopy meets between the rows, combined with post-emergence herbicide treatment	6-8 kg
Sugarcane	Post-transplanting and during vegetative development combined with phytosanitary treatments	6-8 kg
Soybeans	Vegetative development combined with post-emergence herbicide treatment	6-8 kg
Industrial tomato	Post-transplanting, pre-flowering, veraison (change of color), combined with phytosanitary treatments	4-6 kg
Potato	Pre-flowering until tuberification combined with fungicide treatment	4-6 kg
Cotton	Post-emergence and pre- flowering combined with post-emergence/herbicide/fungicide treatments	4-6 kg

COMPOSITION	
Total nitrogen (N)	9%
Nitric nitrogen (N)	9%
Potassium oxide (K <sub>2</sub> O) total	6%
Potassium oxide (K <sub>2</sub> O) soluble in water	6%
Calcium oxide (CaO) total	10%
Calcium oxide (CaO) soluble in water	10%
Boron (B) total	0.10%
Boron (B) soluble in water	0.10%

PHYSICO-CHEMICAL FEATURES	
LIQUID	
pH (sol 1%)	7.2
Conductivity E.C. µS/cm (1‰)	1150
Density (g/cm <sup>3</sup> )/Specific weight	1.50
METHOD OF USE	
	Foliar fertilization

PACKAGING: 6 - 25 KG - COMBO PACK

**PACKAGING: 6, 25 KG - COMBO PACK**

**NOTE:** The activity of SKICC is enhanced when combined with RA.AN L 13186. A COMBO PACK with the 2 products together is now available.

**WARNINGS:** Never mix in the same tank fertilizers containing phosphorus and/or sulfates with fertilizers containing calcium. In presence of irrigation water containing high phosphorus levels it is necessary to add an acidifier before using fertilizers containing calcium.

The product can be mixed with other products exception made for those containing copper, sulfur, mineral oil and emulsions.

It is always advisable to carry out small tests before proceeding with mixing.

\*The choice of the dose is subordinated to various factors and can be varied when necessary. All applications can be repeated in relation to the different crop needs. You can contact our Technical Service for the correct application on specific soils and under specific climate conditions.\*